

Solutions to exercises in Bart Jacobs’s book “Introduction to Coalgebra: Towards Mathematics of States and Observation”

Ryan Tay

some date very far into the future, if ever

a work in progress... draft version 17 September 2025

These are my solutions to all the labelled exercises in [Jacobs \(2017\)](#). This document does not stand on its own; it is meant to supplement the book.

Contents

1	Motivation	2
1.1	Naturalness of Coalgebraic Representations	2
1.2	The Power of Coinduction	4
1.3	Generality of Temporal Logic of Coalgebras	14
1.4	Abstractness of Coalgebraic Notions	17
2	Coalgebras of Polynomial Functors	21
2.1	Constructions on Sets	21
2.2	Polynomial Functors and Their Coalgebras	22
2.3	Final Coalgebras	23
2.4	Algebras	24
2.5	Adjunctions, Cofree Coalgebras, Behaviour-Realisation	25
3	Bisimulations	27
3.1	Relation Lifting, Bisimulations and Congruences	27
3.2	Properties of Bisimulations	27
3.3	Bisimulations as Spans and Cospans	28
3.4	Bisimulations and the Coinduction Proof Principle	28
3.5	Process Semantics	29
	Bibliography and References	30

1 Motivation

1.1 Naturalness of Coalgebraic Representations

Exercise 1.1.1

1. Prove that the composition operation $;$ as defined for coalgebras $S \rightarrow \{\perp\} \cup S$ is associative, i.e. satisfies $s_1 ; (s_2 ; s_3) = (s_1 ; s_2) ; s_3$, for all statements $s_1, s_2, s_3 : S \rightarrow \{\perp\} \cup S$.

Define a statement **skip**: $S \rightarrow \{\perp\} \cup S$ which is a unit for composition $;$ i.e. which satisfies $(\text{skip} ; s) = s = (s ; \text{skip})$, for all $s : S \rightarrow \{\perp\} \cup S$.

2. Do the same for $;$ defined on coalgebras $S \rightarrow \{\perp\} \cup S \cup (S \times E)$.

(In both cases, statements with an associative composition operation and a unit element form a monoid.)

Solution.

1. Recall that the composition operation $;$ was defined as follows:

$$s ; t := \lambda x \in S. \begin{cases} \perp, & \text{if } s(x) = \perp, \\ t(x'), & \text{if } s(x) = x' \in S, \end{cases}$$

for coalgebras $s, t : S \rightarrow \{\perp\} \cup S$. Fix any three coalgebras $s_1, s_2, s_3 : S \rightarrow \{\perp\} \cup S$. Then

$$\begin{aligned} s_1 ; (s_2 ; s_3) &= \lambda x \in S. \begin{cases} \perp, & \text{if } s_1(x) = \perp, \\ (s_2 ; s_3)(x'), & \text{if } s_1(x) = x' \in S, \end{cases} \\ &= \lambda x \in S. \begin{cases} \perp, & \text{if either } s_1(x) = \perp, \text{ or both } s_1(x) = x' \in S \text{ and } s_2(x') = \perp, \\ s_3(x''), & \text{if } s_1(x) = x' \in S \text{ and } s_2(x') = x'' \in S, \end{cases} \\ &= \lambda x \in S. \begin{cases} \perp, & \text{if } (s_1 ; s_2)(x) = \perp, \\ s_3(x''), & \text{if } (s_1 ; s_2)(x) = x'' \in S, \end{cases} \\ &= (s_1 ; s_2) ; s_3. \end{aligned}$$

So the composition operation $;$ is associative.

The coalgebra **skip**: $S \rightarrow \{\perp\} \cup S$ defined by $\text{skip}(x) := x$, for all $x \in S$, satisfies $(\text{skip} ; s) = s = (s ; \text{skip})$ for all coalgebras $s : S \rightarrow \{\perp\} \cup S$.

2. Now we consider the composition operation $;$ defined as follows:

$$s ; t := \lambda x \in S. \begin{cases} \perp, & \text{if } s(x) = \perp, \\ t(x'), & \text{if } s(x) = x' \in S, \\ (x', e), & \text{if } s(x) = (x', e) \in S \times E, \end{cases}$$

for coalgebras $s, t : S \rightarrow \{\perp\} \cup S \cup (S \times E)$. Fix any three coalgebras $s_1, s_2, s_3 : \{\perp\} \cup S \cup (S \times E)$. Then

$$s_1 ; (s_2 ; s_3) = \lambda x \in S. \begin{cases} \perp, & \text{if } s_1(x) = \perp, \\ (s_2 ; s_3)(x'), & \text{if } s_1(x) = x' \in S, \\ (x', e), & \text{if } s_1(x) = (x', e) \in S \times E, \end{cases}$$

$$\begin{aligned}
&= \lambda x \in S. \begin{cases} \perp, & \text{if either } s_1(x) = \perp, \text{ or both } s_1(x) = x' \in S \text{ and } s_2(x') = \perp, \\ s_3(x''), & \text{if } s_1(x) = x' \in S \text{ and } s_2(x') = x'' \in S, \\ (x'', e), & \text{if } s_1(x) = x' \in S \text{ and } s_2(x') = (x'', e) \in S \times E, \\ (x', e), & \text{if } s_1(x) = (x', e) \in S \times E, \end{cases} \\
&= \lambda x \in S. \begin{cases} \perp, & \text{if } (s_1; s_2)(x) = \perp, \\ s_3(x''), & \text{if } (s_1; s_2)(x) = x'' \in S, \\ (x'', e), & \text{if } (s_1; s_2)(x) = (x'', e) \in S \times E, \end{cases} \\
&= (s_1; s_2); s_3.
\end{aligned}$$

So this composition operation $;$ is also associative.

Now define the coalgebra $\text{skip}: S \rightarrow \{\perp\} \cup S \cup (S \times E)$ by $\text{skip}(x) := x$, for all $x \in S$. Then we have $(\text{skip}; s) = s = (s; \text{skip})$ for all coalgebras $s: S \rightarrow \{\perp\} \cup S \cup (S \times E)$. \square

Exercise 1.1.2

Define also a composition monoid $(\text{skip}, ;)$ for coalgebras $S \rightarrow \mathcal{P}(S)$.

Solution. For coalgebras $s, t: S \rightarrow \mathcal{P}(S)$, define

$$s; t := \lambda x \in S. \left(\bigcup_{y \in s(x)} t(y) \right).$$

Then, for coalgebras $s_1, s_2, s_3: S \rightarrow \mathcal{P}(S)$, we have

$$\begin{aligned}
s_1; (s_2; s_3) &= \lambda x \in S. \left(\bigcup_{y \in s_1(x)} (s_2; s_3)(y) \right) \\
&= \lambda x \in S. \left(\bigcup_{y \in s_1(x)} \bigcup_{z \in s_2(y)} s_3(z) \right) \\
&= \lambda x \in S. \left(\bigcup_{z \in (s_1; s_2)(x)} s_3(z) \right) \\
&= (s_1; s_2); s_3.
\end{aligned}$$

Furthermore, defining $\text{skip}: S \rightarrow \mathcal{P}(S)$ by $\text{skip}(x) := \{x\}$ for all $x \in S$, we have

$$\begin{aligned}
(\text{skip}; s) &= \lambda x \in S. \left(\bigcup_{y \in \text{skip}(x)} s(y) \right) \\
&= \lambda x \in S. \left(\bigcup_{y \in \{x\}} s(y) \right) \\
&= \lambda x \in S. s(x) \\
&= s
\end{aligned}$$

and

$$\begin{aligned}
(s; \text{skip}) &= \lambda x \in S. \left(\bigcup_{y \in s(x)} \text{skip}(y) \right) \\
&= \lambda x \in S. \left(\bigcup_{y \in s(x)} \{y\} \right) \\
&= \lambda x \in S. s(x) \\
&= s.
\end{aligned}$$

□

1.2 The Power of Coinduction

Exercise 1.2.1

Compute the `nextdec`-behaviour of $\frac{1}{7} \in [0, 1)$ as in Example 1.2.2.

Solution. We first recall all of the following functions.

1. The final coalgebra `next`: $\{0, \dots, 9\}^\infty \rightarrow \{\perp\} \cup (\{0, \dots, 9\} \times \{0, \dots, 9\}^\infty)$ is defined by

$$\text{next}(\sigma) := \begin{cases} \perp, & \text{if } \sigma \text{ is the empty sequence,} \\ (d, \sigma'), & \text{if } \sigma \text{ has head } d \in \{0, \dots, 9\} \text{ and tail } \sigma' \in \{0, \dots, 9\}^\infty, \text{ i.e. } \sigma = d \cdot \sigma', \end{cases}$$

for all (finite or infinite) sequences $\sigma \in \{0, \dots, 9\}^\infty$.

2. The coalgebra `nextdec`: $[0, 1) \rightarrow \{\perp\} \cup (\{0, \dots, 9\} \times [0, 1))$ is defined by

$$\text{nextdec}(r) := \begin{cases} \perp, & \text{if } r = 0, \\ (d, 10r - d), & \text{if } d \leq 10r < d + 1 \text{ and } d \in \{0, \dots, 9\}, \end{cases}$$

for all $r \in [0, 1)$.

3. The function $\text{beh}_{\text{nextdec}}: [0, 1) \rightarrow \{0, \dots, 9\}^\infty$ is the unique function making

$$\begin{array}{ccc}
\{\perp\} \cup (\{0, \dots, 9\} \times [0, 1)) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_{\{0, \dots, 9\}} \times \text{beh}_{\text{nextdec}})} & \{\perp\} \cup (\{0, \dots, 9\} \times \{0, \dots, 9\}^\infty) \\
\uparrow \text{nextdec} & & \uparrow \cong \text{next} \\
[0, 1) & \xrightarrow{\exists! \text{beh}_{\text{nextdec}}} & \{0, \dots, 9\}^\infty
\end{array}$$

commute.

We wish to compute $\text{beh}_{\text{nextdec}}(\frac{1}{7})$. We see that

$$\begin{aligned}
\text{beh}_{\text{nextdec}}\left(\frac{1}{7}\right) &= \text{next}^{-1} \left(\left(\text{id}_{\{\perp\}} \cup (\text{id}_{\{0, \dots, 9\}} \times \text{beh}_{\text{nextdec}}) \right) \left(\text{nextdec}\left(\frac{1}{7}\right) \right) \right) \\
&= \text{next}^{-1} \left(\left(\text{id}_{\{\perp\}} \cup (\text{id}_{\{0, \dots, 9\}} \times \text{beh}_{\text{nextdec}}) \right) \left(\left(1, \frac{3}{7} \right) \right) \right)
\end{aligned}$$

$$\begin{aligned}
&= \text{next}^{-1} \left(\left(1, \text{beh}_{\text{nextdec}} \left(\frac{3}{7} \right) \right) \right) \\
&= 1 \cdot \text{beh}_{\text{nextdec}} \left(\frac{3}{7} \right).
\end{aligned}$$

Continuing in this fashion,

$$\begin{aligned}
\text{beh}_{\text{nextdec}} \left(\frac{1}{7} \right) &= 1 \cdot \text{beh}_{\text{nextdec}} \left(\frac{3}{7} \right) \\
&= 1 \cdot \left(4 \cdot \text{beh}_{\text{nextdec}} \left(\frac{2}{7} \right) \right) \\
&= 1 \cdot \left(4 \cdot \left(2 \cdot \text{beh}_{\text{nextdec}} \left(\frac{6}{7} \right) \right) \right) \\
&= 1 \cdot \left(4 \cdot \left(2 \cdot \left(8 \cdot \text{beh}_{\text{nextdec}} \left(\frac{4}{7} \right) \right) \right) \right) \\
&= 1 \cdot \left(4 \cdot \left(2 \cdot \left(8 \cdot \left(5 \cdot \text{beh}_{\text{nextdec}} \left(\frac{5}{7} \right) \right) \right) \right) \right) \\
&= 1 \cdot \left(4 \cdot \left(2 \cdot \left(8 \cdot \left(5 \cdot \left(7 \cdot \text{beh}_{\text{nextdec}} \left(\frac{1}{7} \right) \right) \right) \right) \right) \right).
\end{aligned}$$

Therefore $\text{beh}_{\text{nextdec}} \left(\frac{1}{7} \right) = \langle 1, 4, 2, 8, 5, 7, 1, 4, 2, 8, 5, 7, 1, 4, 2, 8, 5, 7, \dots \rangle$.

□

Exercise 1.2.2

Formulate appropriate rules for the function **odds**: $A^\infty \rightarrow A^\infty$ in analogy with the rules (1.7) for **evens**.

Solution. We recall that, for a sequence $\sigma := \langle a_0, a_1, a_2, a_3, \dots \rangle \in A^\infty$, the function **odds** satisfies $\text{odds}(\sigma) = \langle a_1, a_3, a_5, \dots \rangle$, and analogously if σ is a finite sequence. The rules we want **odds** to satisfy are:

$$\frac{\sigma \not\rightarrow}{\text{odds}(\sigma) \not\rightarrow}$$

i.e. **odds** should send the empty sequence to the empty sequence;

$$\frac{\sigma \xrightarrow{a} \sigma' \quad \sigma' \not\rightarrow}{\text{odds}(\sigma) \not\rightarrow}$$

i.e. **odds** should send a singleton sequence $\langle a \rangle$ to the empty sequence; and

$$\frac{\sigma \xrightarrow{a} \sigma' \quad \sigma' \xrightarrow{a'} \sigma''}{\text{odds}(\sigma) \xrightarrow{a'} \text{odds}(\sigma')}$$

i.e. if $\sigma = a \cdot a' \cdot \sigma' \in A^\infty$, where $a, a' \in A$, then $\text{odds}(\sigma) = a' \cdot \text{odds}(\sigma')$.

□

Exercise 1.2.3

Use coinduction to define the empty sequence $\langle \rangle \in A^\infty$ as a map $\{\perp\} \rightarrow A^\infty$.

Fix an element $a \in A$, and similarly define the infinite sequence $\vec{a}: \{\perp\} \rightarrow A^\infty$ consisting of only a s.

Solution. We recall that the final coalgebra $\text{next}: A^\infty \rightarrow \{\perp\} \cup (A \times A^\infty)$ is defined by

$$\text{next}(\sigma) := \begin{cases} \perp, & \text{if } \sigma \text{ is the empty sequence,} \\ (a, \sigma'), & \text{if } \sigma \text{ has head } a \in A \text{ and tail } \sigma' \in A^\infty, \text{ i.e. } \sigma = a \cdot \sigma', \end{cases}$$

for all (finite or infinite) sequences $\sigma \in A^\infty$.

For the coalgebra $\iota_1: \{\perp\} \rightarrow \{\perp\} \cup (A \times \{\perp\})$ defined by $\iota_1(\perp) := \perp$, the unique function $\text{beh}_{\iota_1}: \{\perp\} \rightarrow A^\infty$ making

$$\begin{array}{ccc} \{\perp\} \cup (A \times \{\perp\}) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{beh}_{\iota_1})} & \{\perp\} \cup (A \times A^\infty) \\ \uparrow \iota_1 & & \uparrow \cong \text{next} \\ \{\perp\} & \xrightarrow{\exists! \text{beh}_{\iota_1}} & A^\infty \end{array}$$

commute satisfies $\text{beh}_{\iota_1}(\perp) = \langle \rangle$.

For the coalgebra $c_a: \{\perp\} \rightarrow \{\perp\} \cup (A \times \{\perp\})$ defined by $c_a(\perp) := (a, \perp)$, the unique function $\text{beh}_{c_a}: \{\perp\} \rightarrow A^\infty$ making

$$\begin{array}{ccc} \{\perp\} \cup (A \times \{\perp\}) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{beh}_{c_a})} & \{\perp\} \cup (A \times A^\infty) \\ \uparrow c_a & & \uparrow \cong \text{next} \\ \{\perp\} & \xrightarrow{\exists! \text{beh}_{c_a}} & A^\infty \end{array}$$

commute satisfies $\text{beh}_{c_a}(\perp) = \vec{a} = \langle a, a, a, \dots \rangle$. □

Exercise 1.2.4

Compute the outcome of $\text{merge}(\langle a_0, a_1, a_2 \rangle, \langle b_0, b_1, b_2, b_3 \rangle)$.

Solution. Recall that we defined the coalgebra $m: A^\infty \times A^\infty \rightarrow \{\perp\} \cup (A \times (A^\infty \times A^\infty))$ by

$$m(\sigma, \tau) := \begin{cases} \perp, & \text{if } \sigma \not\rightarrow \text{ and } \tau \not\rightarrow, \\ (a, (\sigma, \tau')), & \text{if } \sigma \not\rightarrow \text{ and } \tau \xrightarrow{a} \tau', \\ (a, (\tau, \sigma')), & \text{if } \sigma \xrightarrow{a} \sigma', \end{cases}$$

for all $\sigma, \tau \in A^\infty$, and that $\text{merge}: A^\infty \times A^\infty \rightarrow A^\infty$ is the unique function making

$$\begin{array}{ccc} \{\perp\} \cup (A \times (A^\infty \times A^\infty)) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{merge})} & \{\perp\} \cup (A \times A^\infty) \\ \uparrow m & & \uparrow \cong \text{next} \\ A^\infty \times A^\infty & \xrightarrow{\exists! \text{merge}} & A^\infty \end{array}$$

commute. Then

$$\text{merge}(\langle a_0, a_1, a_2 \rangle, \langle b_0, b_1, b_2, b_3 \rangle) = \text{next}^{-1} \left((\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{merge})) (m(\langle a_0, a_1, a_2 \rangle, \langle b_0, b_1, b_2, b_3 \rangle)) \right)$$

$$\begin{aligned}
&= \text{next}^{-1} \left((\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{merge})) ((a_0, (\langle b_0, b_1, b_2, b_3 \rangle, \langle a_1, a_2 \rangle))) \right) \\
&= \text{next}^{-1} \left((a_0, \text{merge}(\langle b_0, b_1, b_2, b_3 \rangle, \langle a_1, a_2 \rangle)) \right) \\
&= a_0 \cdot \text{merge}(\langle b_0, b_1, b_2, b_3 \rangle, \langle a_1, a_2 \rangle),
\end{aligned}$$

and so on. Eventually, we obtain $\text{merge}(\langle a_0, a_1, a_2 \rangle, \langle b_0, b_1, b_2, b_3 \rangle) = \langle a_0, b_0, a_1, b_1, a_2, b_2, b_3 \rangle$. \square

Exercise 1.2.5

Is the merge operation associative, i.e. is $\text{merge}(\sigma, \text{merge}(\tau, \rho))$ the same as $\text{merge}(\text{merge}(\sigma, \tau), \rho)$? Give a proof or a counterexample. Is there a neutral element for merge?

Solution. The merge operation is not associative:

$$\begin{aligned}
\text{merge}(\langle a \rangle, \text{merge}(\langle b \rangle, \langle c \rangle)) &= \text{merge}(\langle a \rangle, \langle b, c \rangle) \\
&= \langle a, b, c \rangle,
\end{aligned}$$

whereas

$$\begin{aligned}
\text{merge}(\text{merge}(\langle a \rangle, \langle b \rangle), \langle c \rangle) &= \text{merge}(\langle a, b \rangle, \langle c \rangle) \\
&= \langle a, c, b \rangle,
\end{aligned}$$

for all $a, b, c \in A$.

The neutral element for merge is the empty sequence: for any $\sigma \in A^\infty$, we have $\text{merge}(\sigma, \langle \rangle) = \text{merge}(\langle \rangle, \sigma) = \sigma$. \square

Exercise 1.2.6

Show how to define an alternative merge function which alternately takes two elements from its argument sequences.

Solution. We will define a coalgebra $m_2: A^\infty \times A^\infty \rightarrow \{\perp\} \cup (A \times (A^\infty \times A^\infty))$ so that the desired merge function is the unique function $\text{merge}_2: A^\infty \times A^\infty \rightarrow A^\infty$ making

$$\begin{array}{ccc}
\{\perp\} \cup (A \times (A^\infty \times A^\infty)) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{merge}_2)} & \{\perp\} \cup (A \times A^\infty) \\
\uparrow m_2 & & \uparrow \cong \text{next} \\
A^\infty \times A^\infty & \xrightarrow{\exists! \text{merge}_2} & A^\infty
\end{array}$$

commute. As a motivating example, the desired merge of two infinite streams $\langle a_0, a_1, \dots \rangle$ and $\langle b_0, b_1, \dots \rangle$ should be

$$\text{merge}_2(\langle a_0, a_1, a_2, a_3, \dots \rangle, \langle b_0, b_1, b_2, b_3, \dots \rangle) = \langle a_0, a_1, b_0, b_1, a_2, a_3, b_2, b_3, \dots \rangle.$$

As the diagram above commutes, we would require

$$\text{merge}_2(m_2(\langle a_0, a_1, a_2, a_3, \dots \rangle, \langle b_0, b_1, b_2, b_3, \dots \rangle)) = (a_0, \langle a_1, b_0, b_1, a_2, a_3, b_2, b_3, \dots \rangle)$$

and so m_2 should be defined to satisfy

$$m_2(\langle a_0, a_1, a_2, a_3, \dots \rangle, \langle b_0, b_1, b_2, b_3, \dots \rangle) = (a_0, (\langle a_1, b_0, a_3, b_2, \dots \rangle, \langle b_1, a_2, b_3, a_4, \dots \rangle))$$

Dealing with edge cases separately leads us to the following definition: we define the coalgebra $m_2: A^\infty \times A^\infty \rightarrow \{\perp\} \cup (A \times (A^\infty \times A^\infty))$ as follows.

1. The function m_2 sends the pair $(\langle \rangle, \langle \rangle)$ to \perp , i.e.

$$m_2(\langle \rangle, \langle \rangle) := \perp.$$

2. If $\tau \in A^\infty$ is a non-empty sequence, say $\tau \xrightarrow{a} \tau'$ for some $\tau' \in A^\infty$ and $a \in A$, then

$$m_2(\langle \rangle, \tau) := (a, (\langle \rangle, \tau')).$$

3. If $\sigma = \langle a \rangle$ for some $a \in A$, then

$$m_2(\langle a \rangle, \tau) := (a, (\langle \rangle, \tau))$$

for all $\tau \in A^\infty$.

4. If $\sigma \in A^\infty$ has at least length 2, say $\sigma \xrightarrow{a} \sigma' \xrightarrow{a'} \sigma''$ for some $\sigma', \sigma'' \in A^\infty$ and $a, a' \in A$, then

$$m_2(\sigma, \tau) := \left(a, \left(\text{merge}(\text{odds}(\sigma), \text{evens}(\tau)), \text{merge}(\text{odds}(\tau), \text{evens}(\sigma'')) \right) \right)$$

for all $\tau \in A^\infty$.

Now let $\text{merge}_2: A^\infty \times A^\infty \rightarrow A^\infty$ be the unique function which makes

$$\begin{array}{ccc} \{\perp\} \cup (A \times (A^\infty \times A^\infty)) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{merge}_2)} & \{\perp\} \cup (A \times A^\infty) \\ \uparrow m_2 & & \uparrow \cong \text{next} \\ A^\infty \times A^\infty & \xrightarrow{\exists! \text{merge}_2} & A^\infty \end{array}$$

commute. Fix any $\sigma, \tau \in A^\infty$. We argue by cases on (σ, τ) that this function merge_2 is the desired merge function.

1. If $\sigma = \tau = \langle \rangle$, then $\text{merge}_2(\langle \rangle, \langle \rangle) = \langle \rangle$.
2. If $\sigma = \langle \rangle$ and τ is a non-empty sequence, say $\tau = a \cdot \tau'$ for some $a \in A$ and $\tau' \in A^\infty$, then

$$\text{merge}_2(\langle \rangle, \tau) = a \cdot \text{merge}_2(\langle \rangle, \tau').$$

Thus $\text{merge}_2(\langle \rangle, \tau) = \tau$.

3. If $\sigma = \langle a \rangle$ for some $a \in A$, then

$$\begin{aligned} \text{merge}_2(\langle a \rangle, \tau) &= a \cdot \text{merge}_2(\langle \rangle, \tau) \\ &= a \cdot \tau. \end{aligned}$$

4. If $\sigma = a \cdot a' \cdot \sigma''$ for some $a, a' \in A$ and $\sigma'' \in A^\infty$, then

$$\begin{aligned} \text{merge}_2(\sigma, \tau) &= a \cdot \text{merge}_2\left(\text{merge}(\text{odds}(\sigma), \text{evens}(\tau)), \text{merge}(\text{odds}(\tau), \text{evens}(\sigma''))\right) \\ &= a \cdot \text{merge}_2\left(\text{merge}(a' \cdot \text{odds}(\sigma''), \text{evens}(\tau)), \text{merge}(\text{odds}(\tau), \text{evens}(\sigma''))\right) \\ &= a \cdot a' \cdot \text{merge}_2\left(\text{merge}(\text{odds}(\text{merge}(a' \cdot \text{odds}(\sigma''), \text{evens}(\tau))), \right. \\ &\quad \left. \text{evens}(\text{merge}(\text{odds}(\tau), \text{evens}(\sigma'')))\right), \end{aligned}$$

$$\begin{aligned}
& \text{merge}(\text{odds}(\text{merge}(\text{odds}(\tau), \text{evens}(\sigma''))), \\
& \quad \text{odds}(\text{merge}(\text{evens}(\tau), \text{odds}(\sigma'')))) \\
&= a \cdot a' \cdot \text{merge}_2\left(\text{merge}(\text{evens}(\tau), \text{odds}(\tau)), \text{merge}(\text{evens}(\sigma''), \text{odds}(\sigma''))\right) \\
&= a \cdot a' \cdot \text{merge}_2(\tau, \sigma''),
\end{aligned}$$

as desired. \square

Exercise 1.2.7

1. Define three functions $\text{ex}_i: A^\infty \rightarrow A^\infty$, for $i = 0, 1, 2$, which extract the elements at positions $3n + i$.
2. Define $\text{merge3}: A^\infty \times A^\infty \times A^\infty \rightarrow A^\infty$ satisfying the equation $\text{merge3}(\text{ex}_0(\sigma), \text{ex}_1(\sigma), \text{ex}_2(\sigma)) = \sigma$, for all $\sigma \in A^\infty$.

Solution.

1. Define $c_0, c_1, c_2: A^\infty \rightarrow \{\perp\} \cup (A \times A^\infty)$ as follows:

$$\begin{aligned}
c_0(\sigma) &:= \begin{cases} \perp, & \text{if } \sigma = \langle \rangle, \\ (a, \langle \rangle) & \text{if } \sigma = \langle a \rangle \text{ or } \sigma = \langle a, a' \rangle \text{ for some } a, a' \in A, \\ (a, \sigma'''), & \text{if } \sigma \xrightarrow{a} \sigma' \xrightarrow{a'} \sigma'' \xrightarrow{a''} \sigma''', \end{cases} \\
c_1(\sigma) &:= \begin{cases} \perp, & \text{if } \sigma = \langle \rangle \text{ or } \sigma = \langle a \rangle \text{ for some } a \in A, \\ (a', \langle \rangle) & \text{if } \sigma = \langle a, a' \rangle \text{ for some } a, a' \in A, \\ (a', \sigma'''), & \text{if } \sigma \xrightarrow{a} \sigma' \xrightarrow{a'} \sigma'' \xrightarrow{a''} \sigma''', \end{cases} \\
c_2(\sigma) &:= \begin{cases} \perp, & \text{if } \sigma = \langle \rangle, \text{ or } \sigma = \langle a \rangle, \text{ or } \sigma = \langle a, a' \rangle \text{ for some } a, a' \in A, \\ (a'', \sigma'''), & \text{if } \sigma \xrightarrow{a} \sigma' \xrightarrow{a'} \sigma'' \xrightarrow{a''} \sigma'''. \end{cases}
\end{aligned}$$

Then, for $i \in \{0, 1, 2\}$, the function $\text{ex}_i: A^\infty \rightarrow A^\infty$ is the unique function making

$$\begin{array}{ccc}
\{\perp\} \cup (A \times A^\infty) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{ex}_i)} & \{\perp\} \cup (A \times A^\infty) \\
\uparrow c_i & & \uparrow \cong \text{next} \\
A^\infty & \xrightarrow{\exists! \text{ex}_i} & A^\infty
\end{array}$$

commute.

2. Define the coalgebra $m_3: A^\infty \times A^\infty \times A^\infty \rightarrow \{\perp\} \cup (A \times (A^\infty \times A^\infty \times A^\infty))$ by

$$m_3(\sigma, \tau, \rho) := \begin{cases} \perp, & \text{if } \sigma = \tau = \rho = \langle \rangle, \\ (a, (\langle \rangle, \langle \rangle, \rho')), & \text{if } \sigma = \tau = \langle \rangle \text{ and } \rho \xrightarrow{a} \rho' \text{ for some } a \in A \text{ and } \rho' \in A^\infty, \\ (a, (\langle \rangle, \rho, \tau')), & \text{if } \sigma = \langle \rangle \text{ and } \tau \xrightarrow{a} \tau' \text{ for some } a \in A \text{ and } \tau' \in A^\infty, \\ (a, (\tau, \rho, \sigma')), & \text{if } \sigma \xrightarrow{a} \sigma' \text{ for some } a \in A \text{ and } \sigma' \in A^\infty. \end{cases}$$

Then we let $\text{merge3}: A^\infty \times A^\infty \times A^\infty \rightarrow A^\infty$ be the unique function making

$$\begin{array}{ccc} \{\perp\} \cup (A \times (A^\infty \times A^\infty \times A^\infty)) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{merge3})} & \{\perp\} \cup (A \times A^\infty) \\ \uparrow m_3 & & \uparrow \cong \text{next} \\ A^\infty \times A^\infty \times A^\infty & \xrightarrow{\exists! \text{merge3}} & A^\infty \end{array}$$

commute.

Let us prove that $\text{merge3}(\text{ex}_0(\sigma), \text{ex}_1(\sigma), \text{ex}_2(\sigma)) = \sigma$ for all $\sigma \in A^\infty$, by coinduction. Consider the function $f: A^\infty \rightarrow A^\infty \times A^\infty \times A^\infty$ defined by $f(\sigma) := (\text{ex}_0(\sigma), \text{ex}_1(\sigma), \text{ex}_2(\sigma))$ for all $\sigma \in A^\infty$. We wish to show that $\text{merge3} \circ f = \text{id}_{A^\infty}$.

$$\begin{array}{ccccc} & & \{\perp\} \cup (A \times (A^\infty \times A^\infty \times A^\infty)) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times \text{merge3})} & \{\perp\} \cup (A \times A^\infty) \\ & \nearrow \text{id}_{\{\perp\}} \cup (\text{id}_A \times f) & \uparrow m_3 & & \uparrow \cong \text{next} \\ \{\perp\} \cup (A \times A^\infty) & & & & \\ \uparrow \text{next} \cong & & & & \\ A^\infty & \xrightarrow{f} & A^\infty \times A^\infty \times A^\infty & \xrightarrow{\text{merge3}} & A^\infty \end{array}$$

Let us first show that the left square commutes. It certainly commutes when we chase the empty sequence: $(m_3 \circ f)(\langle \rangle) = \perp = ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times f)) \circ \text{next})(\langle \rangle)$. If $\sigma \in A^\infty$ is a non-empty sequence, say $\sigma \xrightarrow{a} \sigma'$ for some $a \in A$ and $\sigma' \in A^\infty$, then we have

$$\begin{aligned} (m_3 \circ f)(\sigma) &= m_3(\text{ex}_0(\sigma), \text{ex}_1(\sigma), \text{ex}_2(\sigma)) \\ &= (a, (\text{ex}_1(\sigma), \text{ex}_2(\sigma), \text{ex}_0(\sigma'))) \\ &= (a, (\text{ex}_0(\sigma'), \text{ex}_1(\sigma'), \text{ex}_2(\sigma'))) \\ &= ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times f)) \circ \text{next})(\sigma), \end{aligned}$$

where the second-to-last equality can also be proven by coinduction. Therefore the outer square commutes, and so

$$\text{next} \circ (\text{merge3} \circ f) = ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times (\text{merge3} \circ f))) \circ \text{next}.$$

The finality of the coalgebra $\text{next}: A^\infty \rightarrow \{\perp\} \cup (A \times A^\infty)$ now yields $\text{merge3} \circ f = \text{id}_{A^\infty}$. \square

Exercise 1.2.8

Consider the sequential composition function $\text{comp}: A^\infty \times A^\infty \rightarrow A^\infty$ for sequences, described by the three rules:

$$\begin{array}{c} \frac{\sigma \not\rightarrow \quad \tau \not\rightarrow}{\text{comp}(\sigma, \tau) \not\rightarrow} \qquad \frac{\sigma \not\rightarrow \quad \tau \xrightarrow{a} \tau'}{\text{comp}(\sigma, \tau) \xrightarrow{a} \text{comp}(\sigma, \tau')} \\ \frac{\sigma \xrightarrow{a} \sigma'}{\text{comp}(\sigma, \tau) \xrightarrow{a} \text{comp}(\sigma', \tau)} \end{array}.$$

1. Show by coinduction that the empty sequence $\langle \rangle = \text{next}^{-1}(\perp) \in A^\infty$ is a unit element for **comp**, i.e. that $\text{comp}(\langle \rangle, \sigma) = \sigma = \text{comp}(\sigma, \langle \rangle)$.
2. Prove also by coinduction that **comp** is associative, and thus that sequences carry a monoid structure.

Solution.

1. Let $f: A^\infty \rightarrow A^\infty$ be defined by $f(\sigma) := \text{comp}(\langle \rangle, \sigma)$. We will show that the diagram

$$\begin{array}{ccc}
 \{\perp\} \cup (A \times A^\infty) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times f)} & \{\perp\} \cup (A \times A^\infty) \\
 \uparrow \text{next} \cong & & \cong \uparrow \text{next} \\
 A^\infty & \xrightarrow{f} & A^\infty
 \end{array}$$

commutes, which would yield $f = \text{id}_{A^\infty}$ by the finality of the coalgebra **next**.

First, we chase the empty sequence from the bottom left. We see that

$$\begin{aligned}
 (\text{next} \circ f)(\langle \rangle) &= \text{next}(\text{comp}(\langle \rangle, \langle \rangle)) \\
 &= \text{next}(\langle \rangle) \\
 &= \perp,
 \end{aligned}$$

the first rule for **comp**, and

$$\begin{aligned}
 ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times f)) \circ \text{next})(\langle \rangle) &= (\text{id}_{\{\perp\}} \cup (\text{id}_A \times f))(\perp) \\
 &= \perp.
 \end{aligned}$$

Now if $\sigma \in A^\infty$ is a non-empty sequence, say $\sigma \xrightarrow{a} \sigma'$ for some $a \in A$ and $\sigma' \in A^\infty$, we see that

$$\begin{aligned}
 (\text{next} \circ f)(\sigma) &= \text{next}(\text{comp}(\langle \rangle, a \cdot \sigma')) \\
 &= (a, \text{comp}(\langle \rangle, \sigma')) \\
 &= (a, f(\sigma')),
 \end{aligned}$$

by the second rule for **comp** and the definition of f , and

$$\begin{aligned}
 ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times f)) \circ \text{next})(\sigma) &= ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times f))((a, \sigma'))) \\
 &= (a, f(\sigma')).
 \end{aligned}$$

Thus $\text{next} \circ f = (\text{id}_{\{\perp\}} \cup (\text{id}_A \times f)) \circ \text{next}$. This proves that $\text{comp}(\langle \rangle, \sigma) = \sigma$ for all $\sigma \in A^\infty$.

We now show the other equality, that $\text{comp}(\sigma, \langle \rangle) = \sigma$ for all $\sigma \in A^\infty$, we will show that the function $g: A^\infty \rightarrow A^\infty$ defined by $g(\sigma) := \text{comp}(\sigma, \langle \rangle)$ for all $\sigma \in A^\infty$ also satisfies

$$\text{next} \circ g = (\text{id}_{\{\perp\}} \cup (\text{id}_A \times g)) \circ \text{next}.$$

That $(\text{next} \circ g)(\perp) = ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times g)) \circ \text{next})(\perp)$ is the same as with f . Now if $\sigma \in A^\infty$ is such that $\sigma \xrightarrow{a} \sigma'$ for some $a \in A$ and $\sigma' \in A^\infty$, we see that

$$(\text{next} \circ g)(\sigma) = \text{next}(\text{comp}(a \cdot \sigma', \langle \rangle))$$

$$\begin{aligned}
&= (a, \text{comp}(\sigma', \langle \rangle)) \\
&= (a, g(\sigma')),
\end{aligned}$$

by the third rule for **comp** and the definition of g , and

$$\begin{aligned}
((\text{id}_{\{\perp\}} \cup (\text{id}_A \times g)) \circ \text{next})(\sigma) &= ((\text{id}_{\{\perp\}} \cup (\text{id}_A \times g))((a, \sigma'))) \\
&= (a, g(\sigma')).
\end{aligned}$$

Therefore $g = \text{id}_{A^\infty}$, i.e. $\text{comp}(\sigma, \langle \rangle) = \sigma$ for all $\sigma \in A^\infty$.

2. We will define a coalgebra $c: A^\infty \times A^\infty \times A^\infty \rightarrow \{\perp\} \cup (A \times (A^\infty \times A^\infty \times A^\infty))$ such that the functions $h, k: A^\infty \times A^\infty \times A^\infty \rightarrow A^\infty$ given by

$$\begin{aligned}
h(\sigma, \tau, \rho) &:= \text{comp}(\sigma, \text{comp}(\tau, \rho)) \quad \text{and} \\
k(\sigma, \tau, \rho) &:= \text{comp}(\text{comp}(\sigma, \tau), \rho),
\end{aligned}$$

for all $\sigma, \tau, \rho \in A^\infty$, are both coalgebra homomorphisms from c to **next**.

$$\begin{array}{ccc}
& \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times h)} & \\
\{\perp\} \cup (A \times (A^\infty \times A^\infty \times A^\infty)) & & \{\perp\} \cup (A \times A^\infty) \\
& \xleftarrow{\text{id}_{\{\perp\}} \cup (\text{id}_A \times k)} & \\
\uparrow c & & \uparrow \cong \text{next} \\
A^\infty \times A^\infty \times A^\infty & \xrightarrow{h} & A^\infty \\
& \xleftarrow{k} &
\end{array}$$

The finality of **next** would then yield $h = k$.

Define $c: A^\infty \times A^\infty \times A^\infty \rightarrow \{\perp\} \cup (A \times (A^\infty \times A^\infty \times A^\infty))$ by

$$c(\sigma, \tau, \rho) := \begin{cases} \perp, & \text{if } \sigma = \tau = \rho = \langle \rangle, \\ (a, (\langle \rangle, \langle \rangle, \rho')), & \text{if } \sigma = \tau = \langle \rangle \text{ and } \rho = a \cdot \rho' \text{ for some } a \in A \text{ and } \rho' \in A^\infty, \\ (a, (\langle \rangle, \tau', \rho)), & \text{if } \sigma = \langle \rangle \text{ and } \tau = a \cdot \tau' \text{ for some } a \in A \text{ and } \tau' \in A^\infty, \\ (a, (\sigma', \tau, \rho)), & \text{if } \sigma = a \cdot \sigma' \text{ for some } a \in A \text{ and } \sigma' \in A^\infty. \end{cases}$$

Using the rules for **comp**, it is now elementary to check that h and k make their respective diagrams commute. \square

Exercise 1.2.9

Consider two sets A, B with a function $f: A \rightarrow B$ between them. Use finality to define a function $f^\infty: A^\infty \rightarrow B^\infty$ that applies f element-wise. Use uniqueness to show that this mapping $f \mapsto f^\infty$ is ‘functorial’ in the sense that $(\text{id}_A)^\infty = \text{id}_{A^\infty}$ and $(g \circ f)^\infty = g^\infty \circ f^\infty$.

Solution. For a (non-empty) set B , let $\text{next}_B: B^\infty \rightarrow \{\perp\} \cup (B \times B^\infty)$ denote the final coalgebra defined by

$$\text{next}(\sigma) := \begin{cases} \perp, & \text{if } \sigma \text{ is the empty sequence,} \\ (b, \sigma'), & \text{if } \sigma \text{ has head } b \in B \text{ and tail } \sigma' \in B^\infty, \text{ i.e. } \sigma = b \cdot \sigma', \end{cases}$$

for all $\sigma \in B^\infty$. For a function $f: A \rightarrow B$, define a coalgebra $c_f: A^\infty \rightarrow \{\perp\} \cup (B \times A^\infty)$ by

$$c_f(\sigma) := \begin{cases} \perp, & \text{if } \sigma = \langle \rangle, \\ (f(a), \sigma'), & \text{if } \sigma = a \cdot \sigma' \text{ for some } a \in A \text{ and } \sigma' \in A^\infty, \end{cases}$$

for all $\sigma \in A^\infty$. Let $f^\infty: A^\infty \rightarrow B^\infty$ be the unique function making

$$\begin{array}{ccc} \{\perp\} \cup (B \times A^\infty) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_B \times f^\infty)} & \{\perp\} \cup (B \times B^\infty) \\ \uparrow c_f & & \uparrow \cong \text{next}_B \\ A^\infty & \xrightarrow{\exists! f^\infty} & B^\infty \end{array}$$

commute. Then $f(\langle a_0, a_1, a_2, a_3, \dots \rangle) = \langle f(a_0), f(a_1), f(a_2), f(a_3), \dots \rangle$ for all $a_0, a_1, a_2, a_3, \dots \in A$, and analogously for finite sequences.

We see that $c_{\text{id}_A} = \text{next}_A$. So $(\text{id}_A)^\infty = \text{id}_{A^\infty}$ by finality of next_A . Furthermore, for functions $f: A \rightarrow B$ and $g: B \rightarrow C$, we see that

$$\begin{array}{ccc} \{\perp\} \cup (C \times A^\infty) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_C \times f^\infty)} & \{\perp\} \cup (C \times B^\infty) \\ \uparrow c_{g \circ f} & & \uparrow c_g \\ A^\infty & \xrightarrow{f^\infty} & B^\infty \end{array}$$

commutes. Consequently, the outer square in the diagram

$$\begin{array}{ccccc} \{\perp\} \cup (C \times A^\infty) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_C \times f^\infty)} & \{\perp\} \cup (C \times B^\infty) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_C \times g^\infty)} & \{\perp\} \cup (C \times C^\infty) \\ \uparrow c_{g \circ f} & & \uparrow c_g & & \uparrow \cong \text{next}_C \\ A^\infty & \xrightarrow{f^\infty} & B^\infty & \xrightarrow{g^\infty} & C^\infty \end{array}$$

commutes, i.e.

$$\text{next}_C \circ (g^\infty \circ f^\infty) = (\text{id}_{\{\perp\}} \cup (\text{id}_C \times (g^\infty \circ f^\infty))) \circ c_{g \circ f}.$$

The finality of next_C then yields $(g \circ f)^\infty = g^\infty \circ f^\infty$. □

Exercise 1.2.10

Use finality to define a map $\text{st}: A^\infty \times B \rightarrow (A \times B)^\infty$ that maps a sequence $\sigma \in A^\infty$ and an element $b \in B$ to a new sequence in $(A \times B)^\infty$ by adding this b at every position in σ . (This is an example of a ‘strength’ map; see [Exercise 2.5.4](#).)

Solution. Define a coalgebra $c: A^\infty \times B \rightarrow \{\perp\} \cup ((A \times B) \times (A^\infty \times B))$ as follows:

$$c(\sigma, b) := \begin{cases} \perp, & \text{if } \sigma = \langle \rangle, \\ ((a, b), (\sigma', b)), & \text{if } \sigma = a \cdot \sigma' \text{ for some } a \in A \text{ and } \sigma' \in A^\infty, \end{cases}$$

for all $\sigma \in A^\infty$ and $b \in B$. The unique function $\text{st}: A^\infty \times B \rightarrow (A \times B)^\infty$ making

$$\begin{array}{ccc} \{\perp\} \cup ((A \times B) \times (A^\infty \times B)) & \xrightarrow{\text{id}_{\{\perp\}} \cup (\text{id}_{A \times B} \times \text{st})} & \{\perp\} \cup ((A \times B) \times (A \times B)^\infty) \\ \uparrow c & & \uparrow \cong \text{next} \\ A^\infty \times B & \xrightarrow{\exists! \text{st}} & (A \times B)^\infty \end{array}$$

commute will satisfy $\text{st}(\langle a_0, a_1, a_2, \dots \rangle, b) = \langle (a_0, b), (a_1, b), (a_2, b), \dots \rangle$ for all $a_0, a_1, a_2, a_3, \dots \in A$ and $b \in B$, and analogously for finite sequences in A^∞ . \square

1.3 Generality of Temporal Logic of Coalgebras

Exercise 1.3.1

The *nexttime* operator \circ introduced in (1.9) is the so-called **weak** *nexttime*. There is an associated **strong** *nexttime*, given by $\neg \circ \neg$. Note the difference between weak and strong *nexttime* for sequences.

Solution. Recall that, for a sequence coalgebra $c: S \rightarrow \{\perp\} \cup (A \times S)$ and a predicate $P \subseteq S$, we have

$$(\circ P)(x) \quad \text{if and only if} \quad c(x) = \perp \text{ or } c(x) \in A \times P,$$

for all $x \in S$. So,

$$(\circ \neg P)(x) \quad \text{if and only if} \quad c(x) = \perp \text{ or } c(x) \in A \times (S \setminus P),$$

and thus

$$(\neg \circ \neg P)(x) \quad \text{if and only if} \quad c(x) \neq \perp \text{ and } c(x) \notin A \times (S \setminus P).$$

Since the codomain of c is $\{\perp\} \cup (A \times S)$, and since $P \subseteq S$, we can equivalently write this as

$$(\neg \circ \neg P)(x) \quad \text{if and only if} \quad c(x) \in A \times P. \quad \square$$

Exercise 1.3.2

Prove that the ‘truth’ predicate that always holds is a (sequence) invariant. And if P_1 and P_2 are invariants, then so is the intersection $P_1 \cap P_2$. Finally, if P is an invariant, then so is $\circ P$.

Solution. Fix a sequence coalgebra $c: S \rightarrow \{\perp\} \cup (A \times S)$. The truth predicate is the set S itself. Then, for all $x \in S$,

$$(\circ S)(x) \quad \text{if and only if} \quad c(x) = \perp \text{ or } c(x) \in A \times S.$$

Since the codomain of c is $\{\perp\} \cup (A \times S)$, this means that $\circ S = S$, and so S is an invariant.

Now suppose that P_1 and P_2 are invariant, i.e. $P_1 \subseteq \circ P_1$ and $P_2 \subseteq \circ P_2$. Then, for all $x \in S$,

$$\begin{aligned} (\circ(P_1 \cap P_2))(x) & \quad \text{if and only if} \quad c(x) = \perp \text{ or } c(x) \in A \times (P_1 \cap P_2) \\ & \quad \text{if and only if} \quad c(x) = \perp \text{ or } c(x) \in (A \times P_1) \cap (A \times P_2) \\ & \quad \text{if and only if} \quad (c(x) = \perp \text{ or } c(x) \in A \times P_1) \text{ and } (c(x) = \perp \text{ or } c(x) \in A \times P_2) \\ & \quad \text{if and only if} \quad (\circ P_1)(x) \text{ and } (\circ P_2)(x). \end{aligned}$$

Hence $P_1 \cap P_2 \subseteq (\circ P_1) \cap (\circ P_2) = \circ(P_1 \cap P_2)$, and so $P_1 \cap P_2$ is also invariant.

Finally, suppose that P is invariant, i.e. $P \subseteq \circ P$. We aim to show that $\circ P \subseteq \circ \circ P$. Suppose $x \in S$ is such that $(\circ P)(x)$ holds. Then either $c(x) = \perp$ or $c(x) \in A \times P \subseteq A \times \circ P$. Therefore $(\circ \circ P)(x)$ holds. \square

Exercise 1.3.3

1. Show that \Box is an interior operator, i.e. satisfies: $\Box P \subseteq P$, $\Box P \subseteq \Box \Box P$, and $P \subseteq Q \implies \Box P \subseteq \Box Q$.
2. Prove that a predicate P is invariant if and only if $P = \Box P$.

Solution. Fix a sequence coalgebra $c: S \rightarrow \{\perp\} \cup (A \times S)$. Recall that the henceforth operator \Box is defined on predicates $P \subseteq S$ as follows: for all $x \in S$,

$$(\Box P)(x) \text{ if and only if } \text{there exists an invariant } Q \subseteq S \text{ with } x \in Q \subseteq P.$$

In other words, $\Box P$ is the union of all invariants contained in P .

1. If $x \in \Box P$, then there is an invariant $Q \subseteq S$ with $x \in Q \subseteq P$. So $x \in P$ too. Also, Q is an invariant with $x \in Q \subseteq \Box P$. So $x \in \Box \Box P$ as well. Thus $\Box P \subseteq P$ and $\Box P \subseteq \Box \Box P$.

Now suppose $P \subseteq Q \subseteq S$. Then, for any $x \in \Box P$, there is an invariant $R \subseteq S$ with $x \in R \subseteq P \subseteq Q$. So $x \in \Box Q$ as well. Therefore $\Box P \subseteq \Box Q$.

2. For the forward direction, suppose that P is invariant. By definition, $\Box P$ is the union of all invariants contained within P . As P is assumed to be an invariant, we must have $\Box P = P$.

For the converse direction, suppose that $\Box P = P$. We need to show that P is an invariant, i.e. $P \subseteq \circ P$. For any $x \in P = \Box P$, there exists an invariant $Q \subseteq S$ with $x \in Q \subseteq P$. As Q is an invariant, either $c(x) = \perp$ or $c(x) \in A \times Q \subseteq A \times P$. Hence we also have $x \in \circ P$. Therefore $P \subseteq \circ P$, meaning P is an invariant. \square

Exercise 1.3.4

Recall the finite behaviour predicate $\Diamond((-) \nrightarrow)$ from Example 1.3.4.1 and show that it is an invariant: $\Diamond((-) \nrightarrow) \subseteq \circ \Diamond((-) \nrightarrow)$. Hint: For an invariant Q , consider the predicate $Q' = (\neg((-) \nrightarrow) \cap (\circ Q))$.

Solution. Fix a sequence coalgebra $c: S \rightarrow \{\perp\} \cup (A \times S)$. Recall that, for a predicate $P \subseteq S$ and $x \in S$,

$$(\Diamond P)(x) \text{ if and only if } \text{for all invariants } Q \subseteq S, \text{ we have } \neg Q(x) \text{ or } Q \not\subseteq \neg P.$$

That is, $\Diamond P = \neg \Box \neg P$.

Suppose $x \in S$ is such that $\Diamond(x \nrightarrow)$ holds. We need to show that $\circ \Diamond(x \nrightarrow)$ holds, i.e. if $x \xrightarrow{a} x'$ for some $(a, x') \in A \times S$, then $\Diamond(x' \nrightarrow)$ also holds. Fix any invariant $Q \subseteq S$ with $Q \subseteq \neg((-) \nrightarrow)$. We need to show that $\neg Q(x')$.

Following the hint, we consider the predicate

$$Q' := \neg((-) \nrightarrow) \cap (\circ Q).$$

Observe that Q' is an invariant: if $y \in S$ satisfies $Q'(y)$, then there is some $(b, y') \in A \times S$ such that $y \xrightarrow{b} y'$ and $Q(y')$ hold. Then, since $Q \subseteq \neg((-) \nrightarrow)$ and Q is an invariant, we conclude that $Q'(y')$ also holds. So $Q' \subseteq \circ Q'$.

Hence if $Q(x')$ holds, then $Q'(x)$ holds too, contradicting the assumption that $\Diamond(x \nrightarrow)$. \square

Exercise 1.3.5

Let (A, \leq) be a complete lattice, i.e. a poset in which each subset $U \subseteq A$ has a join $\bigvee U \in A$. It is well known that each subset $U \subseteq A$ then also has a meet $\bigwedge U \in A$, given by $\bigwedge U = \bigvee \{a \in A \mid \forall b \in U. a \leq b\}$.

Let $f: A \rightarrow A$ be a monotone function: $a \leq b$ implies $f(a) \leq f(b)$. Recall, e.g. from [Davey and Priestley \(1990, Chapter 4\)](#) that such a monotone f has both a least fixed point $\mu f \in A$ and a greatest fixed point $\nu f \in A$ given by the formulas:

$$\mu f = \bigwedge \{a \in A \mid f(a) \leq a\}, \quad \nu f = \bigvee \{a \in A \mid a \leq f(a)\}.$$

Now let $c: S \rightarrow \{\perp\} \cup (A \times A)$ be an arbitrary sequence coalgebra, with associated nexttime operator \circ .

1. Prove that \circ is a monotone function $\mathcal{P}(S) \rightarrow \mathcal{P}(S)$, i.e. that $P \subseteq Q$ implies $\circ P \subseteq \circ Q$, for all $P, Q \subseteq S$.
2. Check that $\Box P \in \mathcal{P}(S)$ is the greatest fixed point of the function $\mathcal{P}(S) \rightarrow \mathcal{P}(S)$ given by $U \mapsto P \cap \circ U$.
3. Define for $P, Q \subseteq S$ a new predicate $P \mathcal{U} Q \subseteq S$, for ‘ P until Q ’ as the least fixed point of $U \mapsto Q \cup (P \cap \neg \circ \neg U)$. Check that ‘until’ is indeed a good name for $P \mathcal{U} Q$, since it can be described explicitly as

$$\begin{aligned} P \mathcal{U} Q = \{x \in S \mid \exists n \in \mathbb{N}. \exists x_0, x_1, \dots, x_n \in S. \\ x_0 = x \wedge (\forall i < n. \exists a. x_i \xrightarrow{a} x_{i+1}) \wedge Q(x_n) \\ \wedge \forall i < n. P(x_i)\}. \end{aligned}$$

Hint: Don’t use the fixed point definition μ , but first show that this subset is a fixed point, and then that it is contained in an arbitrary fixed point.

(The fixed point definitions that we described above are standard in temporal logic; see e.g. [Emerson \(1990, 3.24–3.25\)](#). The above operation \mathcal{U} is what is called the ‘strong’ until. The ‘weak one’ does not have the negations \neg in its fixed-point description in point 3.)

Solution.

1. For subsets $P, Q \in \mathcal{P}(S)$ with $P \subseteq Q$, and for $x \in S$ such that $(\circ P)(x)$ holds, we have

$$c(x) = \perp \text{ or } c(x) \in A \times P.$$

From the assumption that $P \subseteq Q$, it follows that

$$c(x) = \perp \text{ or } c(x) \in A \times Q,$$

or equivalently, $(\circ Q)(x)$.

2. Fix $P \in \mathcal{P}(S)$ and define $f_P: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ by $f_P(U) := P \cap \circ U$ for all $U \in \mathcal{P}(S)$. Then the greatest fixed point of f_P is

$$\nu(f_P) := \bigcup_{\substack{U \in \mathcal{P}(S), \\ U \subseteq f_P(U)}} U = \bigcup_{\substack{U \in \mathcal{P}(S), \\ U \subseteq P \cap \circ U}} U = \Box P.$$

3. Fix $P, Q \in \mathcal{P}(S)$, and define $f_{P,Q}: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ by

$$f_{P,Q}(U) := Q \cup (P \cap \neg \circ \neg U)$$

for all $U \in \mathcal{P}(S)$. Recall, from [Exercise 1.3.1](#), that

$$\neg \circ \neg U = \{ x \in S : c(x) \in A \times U \}.$$

We wish to show that the set

$$\begin{aligned} U_{P,Q} := Q \cup \Big\{ x \in S : & \text{ there exist } n \in \mathbb{Z}_{>0}, x_0, \dots, x_n \in S \text{ and } a_0, \dots, a_{n-1} \in A \\ & \text{ such that } x = x_0 \xrightarrow{a_0} \dots \xrightarrow{a_{n-1}} x_n \text{ and} \\ & P(x_0), \dots, P(x_{n-1}), \text{ and } Q(x_n) \text{ all hold} \Big\} \end{aligned}$$

is the least fixed point of $f_{P,Q}$.

First, observe that

$$\begin{aligned} f_{P,Q}(U_{P,Q}) &= Q \cup (P \cap \neg \circ \neg U_{P,Q}) \\ &= Q \cup (P \cap \{ x \in S : c(x) \in A \times U_{P,Q} \}) \\ &= Q \cup \{ x \in S : P(x) \text{ and } c(x) \in A \times U_{P,Q} \} \\ &= U_{P,Q}, \end{aligned}$$

so that $U_{P,Q}$ is indeed a fixed point of $f_{P,Q}$.

Now we show that $U_{P,Q}$ is the least fixed point of $f_{P,Q}$. Fix some $B \subseteq S$ with $f_{P,Q}(B) = B$, i.e.

$$Q \cup \{ x \in S : P(x) \text{ and } c(x) \in A \times B \} = B.$$

Then we get $U_{P,Q} \subseteq B$ by induction on the length of finite sequences $x_0, \dots, x_n \in S$ and $a_0, \dots, a_{n-1} \in A$ satisfying $x_0 \xrightarrow{a_0} \dots \xrightarrow{a_{n-1}} x_n$, and $P(x_0) \wedge \dots \wedge P(x_{n-1}) \wedge Q(x_n)$. \square

1.4 Abstractness of Coalgebraic Notions

Exercise 1.4.1

Let $(M, +, 0)$ be a monoid, considered as a category. Check that a functor $F: M \rightarrow \mathbf{Sets}$ can be identified with a **monoid action**: a set X together with a function $\mu: X \times M \rightarrow X$ with $\mu(x, 0) = x$ and $\mu(x, m_1 + m_2) = \mu(\mu(x, m_2), m_1)$.

Solution. Suppose we are given functor $F: M \rightarrow \mathbf{Sets}$. This F sends the unique object $\star \in \mathbf{Obj}(M)$ to a set $F(\star) \in \mathbf{Obj}(\mathbf{Sets})$, and sends each $m \in \mathbf{Arr}(M)$ to a function $Fm: F(\star) \rightarrow F(\star)$. The functoriality of F requires that $F(0) = \text{id}_{F(\star)}$ and $F(m_1 + m_2) = F(m_1) \circ F(m_2)$ for all $m_1, m_2 \in \mathbf{Arr}(M)$. We then define a function $\mu_F: F(\star) \times \mathbf{Arr}(M) \rightarrow F(\star)$ by $\mu_F(x, m) := F(m)(x)$ for all $(x, m) \in F(\star) \times M$.

The equality $\mu_F(x, 0) = x$ for all $x \in F(\star)$ follows the equality $F(0) = \text{id}_{F(\star)}$, while the equality $\mu_F(x, m_1 + m_2) = \mu_F(\mu_F(x, m_2), m_1)$ for all $x \in X$ and $m_1, m_2 \in \mathbf{Arr}(M)$ follows from the equality $F(m_1 + m_2) = F(m_1) \circ F(m_2)$.

Now suppose we are given also given a set X and a function $\mu: X \times \mathbf{Arr}(M) \rightarrow X$ with $\mu(x, 0) = x$ and $\mu(x, m_1 + m_2) = \mu(\mu(x, m_2), m_1)$ for all $x \in X$ and $m, m_1, m_2 \in \mathbf{Arr}(M)$. We then define a functor $F_\mu: M \rightarrow \mathbf{Sets}$ by $F_\mu(\star) := X$, for the unique object $\star \in \mathbf{Obj}(M)$, and $F_\mu(m) := \mu(-, m)$ for each $m \in \mathbf{Arr}(M)$. That F_μ is actually a functor follows from the assumptions on μ .

We then have $F_{\mu_F} = F$ and $\mu_{F_\mu} = \mu$. \square

Exercise 1.4.2

Check in detail that the opposite \mathbb{C}^{op} and the product $\mathbb{C} \times \mathbb{D}$ are indeed categories.

Solution. Let \mathbb{C} and \mathbb{D} be categories.

We defined $\text{Obj}(\mathbb{C}^{\text{op}}) := \text{Obj}(\mathbb{C})$. For $X, Y \in \text{Obj}(\mathbb{C})$, write $\text{hom}_{\mathbb{C}}(X, Y)$ for the set of all morphisms with domain X and codomain Y . We then defined $\text{hom}_{\mathbb{C}^{\text{op}}}(X, Y) := \text{hom}_{\mathbb{C}}(Y, X)$, and we defined a composition $X \xleftarrow{f} Y \xleftarrow{g} Z$ in \mathbb{C}^{op} to be the composition $X \xrightarrow{f} Y \xrightarrow{g} Z$ in \mathbb{C} . The associativity and identity laws for composition in \mathbb{C}^{op} follow from those for \mathbb{C} .

We defined $\text{Obj}(\mathbb{C} \times \mathbb{D}) := \text{Obj}(\mathbb{C}) \times \text{Obj}(\mathbb{D})$. For $X, X' \in \text{Obj}(\mathbb{C})$ and $Y, Y' \in \text{Obj}(\mathbb{D})$, we let $\text{hom}_{\mathbb{C} \times \mathbb{D}}((X, Y), (X', Y')) := \text{hom}_{\mathbb{C}}(X, X') \times \text{hom}_{\mathbb{D}}(Y, Y')$. A composition $(X, Y) \xrightarrow{(f, g)} (X', Y') \xrightarrow{(f', g')} (X'', Y'')$ in $\mathbb{C} \times \mathbb{D}$ is defined to be the composition $(X, Y) \xrightarrow{(f'f, g'g)} (X'', Y'')$. For an object (X, Y) in $\mathbb{C} \times \mathbb{D}$, the identity morphism $\text{id}_{(X, Y)}$ is the pair $(\text{id}_X, \text{id}_Y)$. The associativity and identity laws for composition in $\mathbb{C} \times \mathbb{D}$ follow from those for \mathbb{C} and \mathbb{D} . \square

Exercise 1.4.3

Assume an arbitrary category \mathbb{C} with an object $I \in \mathbb{C}$. We form a new category \mathbb{C}/I , the so-called **slice category** over I , with

objects maps $f: X \rightarrow I$ with codomain I in \mathbb{C}

morphisms from $X \xrightarrow{f} I$ to $Y \xrightarrow{g} I$ are morphisms $h: X \rightarrow Y$ in \mathbb{C} for which the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{h} & Y \\ & \searrow f & \swarrow g \\ & I & \end{array}$$

1. Describe identities and composition in \mathbb{C}/I , and verify that \mathbb{C}/I is a category.
2. Check that taking domains yields a functor $\text{dom}: \mathbb{C}/I \rightarrow \mathbb{C}$.
3. Verify that for $\mathbb{C} = \mathbf{Sets}$, a map $f: X \rightarrow I$ may be identified with an I -indexed family of sets $(X_i)_{i \in I}$, namely where $X_i = f^{-1}(i)$. What do morphisms in \mathbb{C}/I correspond to, in terms of such indexed families?

Solution.

1. The identities and composition in \mathbb{C}/I are simply the identities and composition in \mathbb{C} . So the fact that \mathbb{C}/I is a category follows from \mathbb{C} being a category.
2. We define $\text{dom}: \mathbb{C}/I \rightarrow \mathbb{C}$ as follows: for a morphism h from $X \xrightarrow{f} I$ to $Y \xrightarrow{g} I$ in \mathbb{C}/I , we simply define $\text{dom}(h) := h$. This immediately makes dom a functor from \mathbb{C}/I to \mathbb{C} .
3. The claimed identification is obvious. Now fix a morphism h from $X \xrightarrow{f} I$ to $Y \xrightarrow{g} I$ in \mathbf{Sets}/I , so that the diagram

$$\begin{array}{ccc} X & \xrightarrow{h} & Y \\ & \searrow f & \swarrow g \\ & I & \end{array}$$

in \mathbf{Sets} commutes. This requires that $g(h(x)) = f(x)$ for all $x \in X$. Identifying $X_i := f^{-1}(i)$ and $Y_i := g^{-1}(i)$ for all $i \in I$, we can identify h with a family of functions $(h_i)_{i \in I}$ such that $h_i(x) \in Y_i$ for all $x \in X_i$, for all $i \in I$. \square

Exercise 1.4.4

Recall that for an arbitrary set A we write A^* for the set of finite sequences $\langle a_0, \dots, a_n \rangle$ of elements $a_i \in A$.

1. Check that A^* carries a monoid structure given by concatenation of sequences, with the empty sequence $\langle \rangle$ as a neutral element.
2. Check that the assignment $A \mapsto A^*$ yields a functor $\mathbf{Sets} \rightarrow \mathbf{Mon}$ by mapping a function $f: A \rightarrow B$ between sets to the function $f^*: A^* \rightarrow B^*$ given by $\langle a_0, \dots, a_n \rangle \mapsto \langle f(a_0), \dots, f(a_n) \rangle$. (Be aware of what needs to be checked: f^* must be a monoid homomorphism, and $(-)^*$ must preserve composition of functions and identity functions.)
3. Prove that A^* is the **free monoid on A** : there is the singleton-sequence insertion map $\eta: A \rightarrow A^*$ which is universal among all mappings of A into a monoid. The latter means that for each monoid $(M, 0, +)$ and function $f: A \rightarrow M$ there is a unique monoid homomorphism $g: A^* \rightarrow M$ with $g \circ \eta = f$.

Solution.

1. Concatenation is associative because all the sequences under consideration are finite.
2. That $(-)^*$ preserves composition and identity functions is obvious, so we just check that for a function $f: A \rightarrow B$, the map $f^*: A^* \rightarrow B^*$ is a monoid homomorphism. Fix finite sequences $\langle a_0, \dots, a_n \rangle, \langle a'_0, \dots, a'_k \rangle \in A^*$. Then

$$\begin{aligned} f(\langle a_0, \dots, a_n \rangle \cdot \langle a'_0, \dots, a'_k \rangle) &= f(\langle a_0, \dots, a_n, a'_0, \dots, a'_k \rangle) \\ &= \langle f(a_0), \dots, f(a_n), f(a'_0), \dots, f(a'_k) \rangle \\ &= \langle f(a_0), \dots, f(a_n) \rangle \cdot \langle f(a'_0), \dots, f(a'_k) \rangle \\ &= f(\langle a_0, \dots, a_n \rangle) \cdot \langle a'_0, \dots, a'_k \rangle \end{aligned}$$

and $f(\langle \rangle) = \langle \rangle$. So f^* is a monoid homomorphism.

3. Define $\eta: A \rightarrow A^*$ by $\eta(a) := \langle a \rangle$ for all $a \in A$. Fix a monoid $(M, 0, +)$ and a function $f: A \rightarrow M$. Define $g: A^* \rightarrow M$ by

$$\begin{aligned} g(\langle \rangle) &:= 0 \\ g(\langle a_0, \dots, a_n \rangle) &:= f(a_0) + \dots + f(a_n) \end{aligned}$$

for all $\langle a_0, \dots, a_n \rangle \in A^*$. This g is clearly a monoid homomorphism, using the associativity of $+$ in M . Observe that the diagram

$$\begin{array}{ccc} A & \xrightarrow{\eta} & A^* \\ & \searrow f & \downarrow g \\ & & M \end{array}$$

in \mathbf{Sets} commutes: we have $f(a) = g(\eta(a))$ for all $a \in A$. Now suppose that there is another monoid homomorphism $h: A^* \rightarrow M$ such that the diagram

$$\begin{array}{ccc} A & \xrightarrow{\eta} & A^* \\ & \searrow f & \downarrow h \\ & & M \end{array}$$

in **Sets** commutes. As $h: A^* \rightarrow M$ is a monoid homomorphism and $f = h\eta$, we require that $h(\langle \rangle) = 0$ and

$$\begin{aligned} h(\langle a_0, \dots, a_n \rangle) &= h(\langle a_0 \rangle \cdot \dots \cdot \langle a_n \rangle) \\ &= h(\langle a_0 \rangle) + \dots + h(\langle a_n \rangle) \\ &= h(\eta(a_0)) + \dots + h(\eta(a_n)) \\ &= f(a_0) + \dots + f(a_n) \\ &= g(\langle a_0, \dots, a_n \rangle), \end{aligned}$$

for all $\langle a_0, \dots, a_n \rangle \in A^*$. Therefore $h = g$. \square

Exercise 1.4.5

Recall from (1.3) the statements with exceptions of the form $S \rightarrow \{\perp\} \cup S \cup (S \times E)$.

1. Prove that the assignment $X \mapsto \{\perp\} \cup X \cup (X \times E)$ is functorial, so that the statements are a coalgebra for this functor.
2. Show that all the operations $\text{at}_1, \dots, \text{at}_n, \text{meth}_1, \dots, \text{meth}_m$ of a class as in (1.10) can also be described as a single coalgebra, namely of the functor:

$$X \mapsto D_1 \times \dots \times D_n \times \underbrace{(\{\perp\} \cup X \cup (X \times E)) \times \dots \times (\{\perp\} \cup X \cup (X \times E))}_{m \text{ times}}.$$

Solution.

1. Let $F: \mathbf{Sets} \rightarrow \mathbf{Sets}$ denote this assignment $F(X) := \{\perp\} \cup X \cup (X \times E)$ where all unions are disjoint unions. We define F on morphisms as follows: for functions $f: X \rightarrow Y$, we define $F(f): F(X) \rightarrow F(Y)$ to be the function

$$F(f)(x) := \begin{cases} \perp, & \text{if } x = \perp, \\ f(x), & \text{if } x \in X, \\ (f(x'), e), & \text{if } x = (x', e) \text{ for some } (x', e) \in X \times E. \end{cases}$$

Then $F(\text{id}_X) = \text{id}_{F(X)}$ and $F(gf) = F(g)F(f)$ for all sets X and functions $X \xrightarrow{f} Y \xrightarrow{g} Z$.

2. The functor's definition on morphisms is similar in style with the previous part. \square

Exercise 1.4.6

Recall the nexttime operator \circ for a sequence coalgebra $c: S \rightarrow \mathbf{Seq}(S) = \{\perp\} \cup (A \times S)$ from the previous section. *Exercise 1.3.5.1* says that it forms a monotone function $\mathcal{P}(S) \rightarrow \mathcal{P}(S)$ — with respect to the inclusion order — and thus a functor. Check that invariants are precisely \circ -coalgebras!

Solution. The \circ -coalgebras are simply a subsets $U \subseteq S$ such that $U \subseteq \circ U$. These are precisely what invariants are. \square

2 Coalgebras of Polynomial Functors

2.1 Constructions on Sets

Exercise 2.1.1

Verify in detail the bijective correspondences (2.2), (2.6), (2.11) and (2.16).

Solution. #??

□

Exercise 2.1.2

#??

Solution. #??

□

Exercise 2.1.3

#??

Solution. #??

□

Exercise 2.1.4

#??

Solution. #??

□

Exercise 2.1.5

#??

Solution. #??

□

Exercise 2.1.6

#??

Solution. #??

□

Exercise 2.1.7

#??

Solution. #??

□

Exercise 2.1.8

#??

Solution. #??

□

Exercise 2.1.9

#??

Solution. #??

□

Exercise 2.1.10

#??

Solution. #??

□

Exercise 2.1.11

#??

Solution. #??

□

Exercise 2.1.12

#??

Solution. #??

□

Exercise 2.1.13

#??

Solution. #??

□

Exercise 2.1.14

#??

Solution. #??

□

2.2 Polynomial Functors and Their Coalgebras

Exercise 2.2.1

#??

Solution. #??

□

Exercise 2.2.2

#??

Solution. #??

□

Exercise 2.2.3

#??

Solution. #??

□

Exercise 2.2.4

#??

Solution. #??

□

Exercise 2.2.5

#??

Solution. #??

□

Exercise 2.2.6

#??

Solution. #??

□

Exercise 2.2.7

#??

Solution. #??

□

Exercise 2.2.8

#??

Solution. #??

□

Exercise 2.2.9

#??

Solution. #??

□

Exercise 2.2.10

#??

Solution. #??

□

Exercise 2.2.11

#??

Solution. #??

□

Exercise 2.2.12

#??

Solution. #??

□

2.3 Final Coalgebras

Exercise 2.3.1

#??

Solution. #??

□

Exercise 2.3.2

#??

Solution. #??

□

Exercise 2.3.3

#??

Solution. #??

□

Exercise 2.3.4

#??

Solution. #??

□

Exercise 2.3.5

#??

Solution. #??

□

Exercise 2.3.6

#??

Solution. #??

□

Exercise 2.3.7

#??

Solution. #??

□

Exercise 2.3.8

#??

Solution. #??

□

2.4 Algebras

Exercise 2.4.1

#??

Solution. #??

□

Exercise 2.4.2

#??

Solution. #??

□

Exercise 2.4.3

#??

Solution. #??

□

Exercise 2.4.4

#??

Solution. #??

□

Exercise 2.4.5

#??

Solution. #??

□

Exercise 2.4.6

#??

Solution. #??

□

Exercise 2.4.7

#??

Solution. #??

□

Exercise 2.4.8

#??

Solution. #??

□

Exercise 2.4.9

#??

Solution. #??

□

Exercise 2.4.10

#??

Solution. #??

□

2.5 Adjunctions, Cofree Coalgebras, Behaviour-Realisation

Exercise 2.5.1

#??

Solution. #??

□

Exercise 2.5.2

#??

Solution. #??

□

Exercise 2.5.3

#??

Solution. #??

□

Exercise 2.5.4

This exercise describes ‘strength’ for endofunctors on **Sets**. In general, this is a useful notion in the theory of datatypes (*Cockett and Spencer, 1992*), (*Cockett and Spencer, 1995*) and of computations (*Moggi, 1991*); see Section 5.2 for a systemic description.

Let $F: \mathbf{Sets} \rightarrow \mathbf{Sets}$ be an arbitrary functor. Consider for sets X, Y the strength map

$$F(X) \times Y \xrightarrow{\text{st}_{X,Y}} F(X \times Y)$$

$$(u, y) \longmapsto F(\lambda x \in X. (x, y))(u)$$

1. Prove that this yields a natural transformation $F(-) \times (-) \xRightarrow{\text{st}} F((-) \times (-))$, where both the domain and codomain are functors $\mathbf{Sets} \times \mathbf{Sets} \rightarrow \mathbf{Sets}$.
2. Describe this strength map for the list functor $(-)^*$ and for the powerset functor \mathcal{P} .

Solution. #??

□

Exercise 2.5.5

#??

Solution. #??

□

Exercise 2.5.6

#??

Solution. #??

□

Exercise 2.5.7

#??

Solution. #??

□

Exercise 2.5.8

#??

Solution. #??

□

Exercise 2.5.9

#??

Solution. #??



Exercise 2.5.10

#??

Solution. #??



Exercise 2.5.11

#??

Solution. #??



Exercise 2.5.12

#??

Solution. #??



Exercise 2.5.13

#??

Solution. #??



Exercise 2.5.14

#??

Solution. #??



Exercise 2.5.15

#??

Solution. #??



Exercise 2.5.16

#??

Solution. #??



Exercise 2.5.17

#??

Solution. #??



3 Bisimulations

3.1 Relation Lifting, Bisimulations and Congruences

Exercise 3.1.1

#??

Solution. #??

□

Exercise 3.1.2

#??

Solution. #??

□

Exercise 3.1.3

#??

Solution. #??

□

Exercise 3.1.4

#??

Solution. #??

□

Exercise 3.1.5

#??

Solution. #??

□

Exercise 3.1.6

#??

Solution. #??

□

3.2 Properties of Bisimulations

Exercise 3.2.1

#??

Solution. #??

□

Exercise 3.2.2

#??

Solution. #??

□

Exercise 3.2.3

#??

Solution. #??

□

Exercise 3.2.4

#??

Solution. #??

□

Exercise 3.2.5

#??

Solution. #??

□

Exercise 3.2.6

#??

Solution. #??

□

Exercise 3.2.7

#??

Solution. #??

□

3.3 Bisimulations as Spans and Cospans**Exercise 3.3.1**

#??

Solution. #??

□

Exercise 3.3.2

#??

Solution. #??

□

Exercise 3.3.3

#??

Solution. #??

□

Exercise 3.3.4

#??

Solution. #??

□

3.4 Bisimulations and the Coinduction Proof Principle**Exercise 3.4.1**

#??

Solution. #??

□

Exercise 3.4.2

#??

Solution. #??

□

Exercise 3.4.3

#??

Solution. #??

□

Exercise 3.4.4

#??

Solution. #??



Exercise 3.4.5

#??

Solution. #??



Exercise 3.4.6

#??

Solution. #??



Exercise 3.4.7

#??

Solution. #??



3.5 Process Semantics

Exercise 3.5.1

#??

Solution. #??



Exercise 3.5.2

#??

Solution. #??



Exercise 3.5.3

#??

Solution. #??



Exercise 3.5.4

#??

Solution. #??



Bibliography and References

- J. Robin B. Cockett and Dwight Spencer. Strong categorical datatypes I. In Robert A. G. Seely, editor, *International Meeting on Category Theory 1991*, volume 13, pages 141–169. Canadian Mathematical Society Proceedings, AMS, Montreal, 1992.
- J. Robin B. Cockett and Dwight Spencer. Strong categorical datatypes II: A term logic for categorical programming. *Theoretical Computer Science*, 139:69–113, 1995.
DOI: [https://doi.org/10.1016/0304-3975\(94\)00099-5](https://doi.org/10.1016/0304-3975(94)00099-5).
- Brian A. Davey and Hilary A. Priestley. *Introduction to Lattices and Order*. Cambridge University Press, 1990.
DOI: <https://doi.org/10.1017/CB09780511809088>.
- E. Allen Emerson. Temporal and modal logic. In Jan van Leeuwen, editor, *Handbook of Theoretical Computer Science*, volume B: Formal Models and Semantics, pages 995–1072. Elsevier B.V., 1990.
DOI: <https://doi.org/10.1016/B978-0-444-88074-1.50021-4>.
- Bart Jacobs. *Introduction to Coalgebra: Towards Mathematics of States and Observation*. Cambridge University Press, 2017.
DOI: <https://doi.org/10.1017/CB09781316823187>.
- Eugenio Moggi. Notions of computation and monads. *Information and Computation*, 93(1):55–92, 1991.
DOI: [https://doi.org/10.1016/0890-5401\(91\)90052-4](https://doi.org/10.1016/0890-5401(91)90052-4).